

First Record of the Genus *Janolus* Bergh, 1884 (Opisthobranchia: Arminacea: Zephyrinidae) from the Pacific Coast of South America, with the Description of a New Species

by

MARIA ANGELICA FISCHER

Departamento de Biología de Organismos y Sistemas, Laboratorio de Zoología,
Facultad de Biología, Universidad de Oviedo, 33071 Oviedo, Asturias, Spain

JUAN LUCAS CERVERA

Departamento de Biología Animal, Vegetal y Ecología, Facultad de Ciencias del Mar;
Universidad de Cádiz, Apdo. 40, 11510 Puerto Real (Cádiz), Spain

AND

JESÚS ORTEA

Departamento de Biología de Organismos y Sistemas, Laboratorio de Zoología,
Facultad de Biología, Universidad de Oviedo, 33071 Oviedo, Asturias, Spain

Abstract. A new species of the nudibranch genus *Janolus* Bergh, 1884, from the Chilean coast is described. The external and internal anatomy and coloration are described. The species, *Janolus chilensis* sp. nov., is compared with those most similar to it. *Janolus chilensis* constitutes the first record of the genus on the Chilean and South American Pacific coasts.

INTRODUCTION

The genus *Janolus* Bergh, 1884, includes only two known species from the American Pacific coast: *J. barbarensis* (Cooper, 1863) and *J. fuscus* (O'Donoghue, 1924). Behrens (1991) illustrated a third undescribed species identified as *Janolus* sp. 1. The known geographical range for these species extends more or less along the North American Pacific coast, depending on the species, but never south of the Gulf of California (Behrens, 1991).

On the Chilean coasts the suborder Arminacea is represented by only one species, *Armina cuvieri* (d'Orbigny, 1837). In this paper we describe a new species of zephyrinid nudibranch, recently illustrated and identified as *Janolus* sp. 1 by Schrödl (1996: tab. 6, fig. 34), from two specimens collected on the north Chilean coast.

SYSTEMATIC DESCRIPTION

Suborder ARMINACEA Odhner, 1934

Family ZEPHYRINIDAE Iredale & O'Donoghue,
1923

Genus *Janolus* Bergh, 1884

Janolus chilensis Fischer, Cervera & Ortea,
sp. nov.

(Figures 1–5)

Material: Holotype: One specimen, 45 mm in length, collected on hydrozoan and bryozoan colonies attached to floats and culture lines substrates, at 3 m depth, Iquique (20°20'S; 70°05'W), north coast of Chile, 10 August 1993,




Figure 1

Janolus chilensis Fischer, Cervera & Ortea, sp. nov. A. Dorsal view of the living animal. B. Color pattern of a ceras. C. Detail of the anal papilla. D. Ventral view of the anterior part of the animal. Key: ap, anal papilla; bl, blue; cr, caruncle; dgl, digestive gland; mo, mouth; or, orange; ot, oral tentacle; ow, opaque white; rh, rhinophore.

M. Angelica Fischer coll. This specimen, which is not dissected, has been deposited in the collections of the Museo Nacional de Historia Natural (MNHN) de Santiago de Chile, catalogue number 201621.

Paratype: One specimen dissected, 40 mm in length, collected concurrently with the holotype, has been also deposited in the collections of the MNHN, catalogue number 201622.




Figure 2

J. chilensis Fischer, Cervera & Ortea, sp. nov. Several cerata showing branching of digestive gland.

Description: Body stout, broadest anteriorly, narrower behind rhinophores, and ending in long, fine tail. Body of live animal translucent white; rhinophores also translucent with blue apex (Figure 1A). Cerata translucent with opaque white apical band, orange subapical band, and narrow translucent band between them (Figure 1B). Branches of digestive gland in cerata translucent brown. Tail translucent with thin central opaque white line. Head wider than rest of body. Pair of short, pointed oral tentacles present in front of head, near mouth. Edge of foot smooth and anteriorly rounded with deep transverse groove (Figure 1D). Rhinophores perfoliate, with 16 thin complete or incomplete transverse lamellae, shortest at base of club. Between rhinophores is an obvious orange bilobed caruncle formed by many lumpy groups (Figure 1A). Anterior to it is also an orange spot. Anus at end of body, where tail begins; anus a short tube opening like flower with nine folds (Figure 1A, C). No anal gland observed. Cerata arranged dorsoventrally around body, 36 on each side. At head, are two alternating ceratal rows. In middle of body are three alternating ceratal rows. Behind anus, cerata form only single row. Cerata shorter and thinner anteriorly and longer and thicker in central body region. Cerata smooth, digitiform, inflated in apical zone with little nipple-shaped end. Digestive gland in cerata appears as slender mostly unbranched tributary, but some have two to five branches that terminate in subapical zone (Figure 2).

Jaws strong, roughly triangular in shape, each side with six broad teeth (Figure 3A, B). Radula broad and well developed. Radular formula of 40 mm specimen $18 \times 18-24.1.18-24$. Rachidian tooth smooth, straight, large, winged, with two small denticles on each side (Figure 4A, B,r). Lateral teeth sickle-shaped. Cusp of lateral teeth becomes shorter from inner portion toward outer edge of radula. Three inner lateral teeth possess two to four denticles (Figure 4A, B).

Reproductive system diaulic (Figure 5A). Penis smooth, muscular, broad, and elongate with pointed end (Figure 5B). Penial sac thick and muscular. Vas deferens narrow, relatively short, and coiled twice, without differentiated prostatic portion. Junction of vas deferens and



Figure 3

J. chilensis Fischer, Cervera & Ortea, sp. nov. Paratype, MNHN 201622. A, Jaws (scanning electron micrograph). B, Detail of the denticles of the jaws (scanning electron micrograph).


ampulla inside female gland. Ampulla thin, elongate, and refringent, continuing distally as slender hermaphroditic duct, which branches into three ducts. Female gland massive, surrounding bursa copulatrix. Bursa rounded joining female gland via relatively long vagina. Both bursa copulatrix and vagina represented by discontinuous line in figure 5A, according to their arrangement.

DISCUSSION

Janolus chilensis can be separated from the other two American Pacific *Janolus* species (see Table 1). Thus, *J. barbarendis* differs from our species by having an anal gland and a receptaculum seminis. It also lacks lateral

Table 1
Comparative morphology of *Janolus chilensis* sp. nov. and its most similar Pacific species.

Species	Color	Digestive ceratal branches	Anal gland	Jaws	Radular formula and teeth	Receptaculum seminis	Bursa copulatrix	Penis	References
<i>Janolus barbarensis</i> (Cooper, 1863)	body translucent white; cerata with gold sub- apical ring and blue tips, branches of di- gestive gland brown; rhino- phores with lemon yellow subapical ring and blue tip; or- ange caruncle	branched	present	7-9 denticles	16 × 27.1.27; ra- chidian narrow, without denti- cles; inner later- al teeth about 6 denticles; remaining later- al smooth	spherical, thin- walled, dis- tal	short, spheri- cal, serial, proximal	large, muscu- lar, thickest near the middle	Cockerell & Eliot (1905); MacFarland (1966); Gos- liner (1981, 1982)
<i>J. fuscus</i> O'Dono- ghue, 1924	body translucent white, with mid- dorsal red- brown lines; cerata with sub- apical yellow and apical opaque white bands, branches of the digestive gland brown; rhinophores pink and opaque white tip; red- brown caruncle	unbranched	absent	10-13 denticles	21-26 × 22- 25.1.22-25; ra- chidian broad, denticulate; inner laterals 2 denticles; re- maining laterals smooth	pyriform, dis- tal	elongate, semi-serial, proximal	conical, thickened posteriorly	O'Donoghue (1924); MacFarland (1966); Gos- liner (1981, 1982)
<i>J. chilensis</i> sp. nov.	body translucent white; cerata with opaque white tip and orange subapi- cal ring; branch- es of the diges- tive gland pale brown; rhino- phores tip dark blue; or- ange caruncle	branched	absent	6-7 denticles	18 × 18-24.1.18- 24; rachidian narrow, denticu- late; inner three lateral with 2-4 denticles; re- maining lateral smooth	absent	spherical, semi-serial	thick, muscu- lar, conical, pointed	present study



A




Figure 4

J. chilensis Fischer, Cervera & Ortea, sp. nov. Paratype, MNHN 201622. A. Schematic view of the radular teeth in a half-row. B. Scanning micrograph of several half-rows of the radula. Key: il, inner lateral teeth; ol, outer lateral teeth; r, rachidian tooth.

denticles on the rachidian tooth. On the other hand, *J. fuscus* also has a receptaculum seminis. In the genus *Janolus*, the presence of one or two receptacles varies intraspecifically. Following the revisions of Gosliner (1981, 1982) and Miller & Willan (1986), the species recognized as having only a bursa copulatrix are *J. conis* Marcus, 1955 (has a bursa/vagina), *J. mucloc* (Marcus, 1958), *J. toyamensis* Baba & Abe, 1970, and *J. eximius* Miller & Willan, 1986; and the species recognized as having only a receptaculum seminis is *J. hyalinus* (Alder & Hancock, 1854). There are also other species that have both types of receptacles: *J. cristatus* (Delle Chiaje, 1841), *J. barbarensis* (Cooper, 1863), *J. novozealandicus* (Eliot, 1907), *J. fuscus* O'Donoghue, 1924, *J. capensis* Bergh, 1907, *J. longidentatus* Gosliner, 1981, *J. ignis* Miller & Willan, 1986 (has a bursa/vagina), and *J. mokohinau* Mil-

ler & Willan, 1986 (has a bursa/vagina). However, there are several species whose reproductive systems remain unknown: *J. australis* Bergh, 1884, *J. indicus* (Eliot, 1909), *J. mirabilis* Baba & Abe, 1970, *J. preclarus* (Bouchet, 1975). After the Gosliner and Miller & Willan papers, Ortea & Llera (1988) described *J. faustoi* from the Canary Islands, but with no details of the reproductive system. A bursa copulatrix surrounded by the female gland was also described in *J. mucloc* by Marcus (1958), similar to *J. chilensis*.

On the other hand, the coloration in some representatives of the Zephyrinidae is variable, like *Bonisa nakaza* Gosliner, 1981. This species is recorded with three different patterns of coloration, but in *Janolus* this variability has not been described. *J. chilensis* has a distinct pattern of coloration. It is clearly different from the other




Figure 5

J. chilensis Fischer, Cervera & Ortea, sp. nov. Paratype, MNHN 201622. A. Reproductive system. B. Detail of the penis. Key: am, ampulla; bc, bursa copulatrix (discontinuous line), fgl, female gland; ps, penial sheath; va, vagina (discontinuous line); vd, vas deferens.

species of *Janolus* from the American Pacific. *Janolus fuscus* has mid-dorsal lines of red-brown, yellow and white cerata, and pink and opaque white rhinophores. *Janolus barbarensis* has cerata with a subapical gold ring, and rhinophores with a subapical lemon-yellow ring and blue tip. *J. chilensis* has opaque white ceratal tips, a subapical orange ring, and the tips of the rhinophores are blue (Table 1). The coloration of the remaining species

of *Janolus* is clearly different. The rachidian teeth of *J. chilensis* are similar to the rachidian teeth of *J. barbarensis*. The two inner lateral teeth of *J. barbarensis* are denticulate, whereas in *J. chilensis* the three inner teeth are denticulate.

Etymology: The specific name, *chilensis*, is from Chile, where the new species was collected.

ACKNOWLEDGMENTS

We are grateful to the University Arturo Pratt in Iquique (Chile) for giving us valuable assistance in collecting specimens, and the Sarmentia Marine Cultures for allowing access to their floating docks; the Electron Microscopy Service of the Faculty of Medicine of the University of Oviedo (Spain); and especially for financial support from the Instituto de Cooperación Iberoamericana (ICI) to M.A.F. Dr. Terrence M. Gosliner (California Academy of Sciences, San Francisco, California) critically read the manuscript and provided valuable comments.

LITERATURE CITED

- BEHRENS, D. W. 1991. Pacific Coast Nudibranchs. 2nd ed. Sea Challengers: Monterey, California. 107 pp.
- COCKERELL, T. & C. ELIOT. 1905. Notes on a collection of Californian nudibranchs. *Journal of Malacology* 12(3):31–53.
- COOPER, J. 1863. On new or rare Mollusca inhabiting the coast of California. 2. *Proceedings of the California Academy of Sciences* 3:56–60.
- GOSLINER, T. M. 1981. The South African Janolidae (Mollusca, Nudibranchia) with the description of a new genus and two new species. *Annals of the South African Museum* 86(1):1–42.
- GOSLINER, T. M. 1982. The genus *Janolus* (Nudibranchia: Arminacea) from the Pacific Coast of North America, with a reinstatement of *Janolus fuscus* O'Donoghue, 1924. *The Veliger* 24(3):219–226.
- MACFARLAND, F. 1966. Studies of opisthobranchiate mollusks of the Pacific coast of North America. *Memoirs of the California Academy of Sciences* 6:1–546.
- MARCUS, E. 1958. On western Atlantic opisthobranchiate gastropods. *American Museum Novitates* 1906:1–82.
- MILLER, M. C. & R. C. WILLAN. 1986. A review of the New Zealand arminacean nudibranchs (Opisthobranchia: Arminacea). *New Zealand Journal of Zoology* 13:337–408.
- O'DONOGHUE, C. H. 1924. Notes on the nudibranchiate Mollusca from the Vancouver Island Region. 4. *Transactions of the Royal Canadian Institute* 15(1):1–33.
- ORTEA, J. A. & E. M. LLERA. 1988. Una nueva especie de *Janolus* Bergh, 1884 (Mollusca: Nudibranchia) dedicada a la memoria de Fausto González. *Publicações Ocasionais da Sociedade Portuguesa de Malacologia* 11:33–38.
- SCHRÖDL, M. 1996. Nudibranchia y Sacoglossa de Chile: distribución, descripción externa y clave. *Gayana Zoología* 60(1):17–62.